

Table 6. Spectral fitting results for NGC3690.

Parameter	Power-law	Single Temperature Raymond-Smith	Double Temperature Raymond Smith	Raymond Smith + Power-law	Triple temperature Raymond Smith	cev
kT (KeV)		$4.25^{+0.36}_{-0.45}$	$0.83^{+0.03}_{-0.03}$ $10.3^{+5.9}_{-2.4}$	$0.83^{+0.02}_{-0.04}$	$0.285^{+0.187}_{-0.063}$ $0.871^{+0.427}_{-0.676}$ $11.17^{+6.96}_{-3.13}$	57.08
Γ	$2.23^{+0.12}_{-0.12}$			$1.56^{+0.11}_{-0.11}$		
$\alpha^\dagger$						0.22
$N_H (10^{20} \text{ cm}^{-2})$	$5.50^{+0.7}_{-0.7}$	$2.11^{+0.38}_{-0.33}$	$1.60^{+0.42}_{-0.40}$	$2.42^{+0.63}_{-0.46}$	$2.031^{+0.283}_{-0.472}$	3.21
χ^2 / d.o.f.	490.0/242	624.5/242	287.8/235	291.5/235	290.3/237	309.
Flux ‡ (0.1-2.0keV)	9.72	8.85	8.5	9.0	8.95	8
Flux ‡ (2.0-10.0keV)	7.68	9.59	10.8	10.8	10.3	9
Total Luminosity* (0.1-10.0keV)	8.3	8.8	9.2	9.46	9.2	

† The slope of power-law distribution of temperatures in the cevmekl model.

‡ The fluxes are in units of $10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$.

* The luminosity is in units of $10^{41} \text{ erg s}^{-1}$.

ROSAT and ASCA Observations of X-ray Luminous starburst Galaxies : NGC3310 and NGC3690

A.L. Zezas¹, I. Georgantopoulos² and M. J. Ward¹

¹ *Department of Physics and Astronomy, University of Leicester, Leicester, LE1 7RH*

² *National Observatory of Athens, Lofos Koufou, Palaia Penteli, 15236, Athens, Greece*

1 February 2008

ABSTRACT

We present ROSAT (HRI and PSPC) and ASCA observations of the two luminous ($L_x \sim 10^{41-42}$ erg s⁻¹) star-forming galaxies NGC3310 and NGC3690. The HRI shows clearly that the sources are extended with the X-ray emission in NGC3690 coming from at least three regions. The combined 0.1-10 keV spectrum of NGC3310 can be described by two components, a Raymond-Smith plasma with temperature $kT = 0.81^{+0.09}_{-0.12}$ keV and a hard power-law, $\Gamma = 1.44^{+0.20}_{-0.11}$, (or alternatively a harder Raymond-Smith plasma with $kT \sim 15$ keV), while there is no substantial excess absorption above the Galactic. The soft component emission is probably due to a super-wind while the nature of the hard emission is more uncertain with likely origins, X-ray binaries, inverse Compton scattering of IR photons, an AGN or a very hot gas component ($\sim 10^8$ K). The spectrum of NGC3690 is similar, with $kT = 0.83^{+0.02}_{-0.04}$ keV and $\Gamma = 1.56^{+0.11}_{-0.11}$. We also employ more complicated models such as a multi-temperature thermal plasma, a non-equilibrium ionization code or the addition of a third softer component which improve the fit but not at a statistically significant level ($< 2\sigma$). These results are similar to recent results on the archetypal star-forming galaxies M82 and NGC253.

Key words: galaxies: starburst-galaxies: merger-X-rays: galaxies-galaxies: individual: NGC3690-galaxies: individual: NGC3310

1 INTRODUCTION

Significant X-ray emission from star-forming galaxies has been known since the *EINSTEIN* epoch (eg Stewart et al. 1982). Their X-ray spectrum in the 0.3-3.5 keV band can be fitted by a thermal plasma of temperature about 2 keV with an absorption often higher than the Galactic value (eg. Fabbiano 1988, Kim et al. 1992). However, simple spectral fits suggested that the emission mechanism may be more complicated than a single temperature thermal plasma. Indeed, for a few nearby star-forming galaxies (eg M82 and NGC253) several discrete sources, mostly X-ray binaries, are also resolved (Long & Van Speybroeck 1983). The *EINSTEIN* HRI provided the first evidence for extended X-ray features (eg Watson et al. 1984, Fabbiano and Trinchieri 1984).

The effective area of *ROSAT* PSPC provided the opportunity to dramatically increase the number of star-forming galaxies studied in X-rays. Hot gas halos and outflows, have also been observed, around many star-forming galaxies (eg. Read, Ponman & Wolstencroft 1995, Della Ceca, Griffiths & Heckman 1996, Read, Ponman & Strickland 1997), as was anticipated based on previous optical (Heckman et al. 1990) and theoretical work (e.g. MacLow and Mc Cray

1988). This wind can be produced by the supply of mechanical energy via stellar winds from evolved massive stars and numerous supernovae. The hot gas is accelerated outwards forming an expanding superbubble, which on reaching the extent of the minor axis of the galaxy, blows out following the onset of the Rayleigh-Taylor instability. These outflows can extend for several kpc along the minor axis of the galaxy, and have major influence on its subsequent evolution. This is especially so in the case of dwarf galaxies, where because of their weak gravitational potential, the wind can remove the interstellar medium causing star formation to cease, (eg Dekel & Silk 1986, Heckman et al. 1995). Combined ROSAT and ASCA observations of star-forming galaxies, (eg Serlemitsos, Ptak & Yaqoob 1996) suggest that the soft X-ray band can be fitted with a thermal plasma and a power-law model. The plasma temperature is $\sim 0.6 - 0.8$ keV, with an absorbing column $N_H > 10^{21}$ cm⁻², well above the Galactic value, and the power-law photon index is ~ 1.7 . The hard X-ray band can be fitted equally well with a high temperature thermal plasma model, $kT > 6$ keV, (cf. Ohashi et al. 1990). The soft emission may emanate from the super-wind while the origin of the harder power-law X-ray emission may be due to either X-ray binaries or Inverse Compton scattering

of infrared-photons by the relativistic electrons generated by the supernovae (eg Rieke et al. 1980). The presence of very hot gas ($\sim 10^8\text{K}$), or a weak active nucleus cannot be ruled out. However, medium spectral resolution, wide energy band (0.5-10 keV) observations of NGC253 and M82 with ASCA, show that the above two component models do not give an acceptable description of the data (Ptak et al 1997, Moran & Lehnert 1997). Even the introduction of a third thermal component ($kT \sim 0.3\text{keV}$) still provides a poor fit to the high-resolution ASCA SIS spectrum (Moran & Lehnert 1997). This suggests that the emission mechanisms are more complex than these simple models can explain. Since the X-ray spectra of Galactic supernova remnants (eg Hughes & Singh 1994) cannot be well-fit by a simple Raymond-Smith model (their emission is dominated by non-equilibrium processes) it is likely that we are dealing with a similar situation in starforming galaxies.

One of the interesting results of the *ROSAT* All-Sky Survey was the discovery of star-forming galaxies with X-ray luminosities $\sim 10^{42}\text{ergs}^{-1}$, approaching those of Seyfert galaxies (eg Moran et al 1995). In order to gain further insight into the nature of this powerful X-ray emission and to compare starburst properties with those of the archetypal (less luminous) star-forming galaxies, we have cross-correlated the *ROSAT* All Sky Survey Bright Source Catalogue, RASSBSC, (Voges et al. 1996) with the spectral atlas of nearby galaxies of Ho, Filippenko and Sargent (1995). The advantage of the latter sample is that it has high quality optical spectra and thus provides us with a reliable classification for each galaxy. The sample of Ho et al. contains galaxies with declination $\delta > 0^\circ$ and magnitude $B < 12.5$ while the RASS contains almost 20,000 sources over the whole sky. The cross-correlation yielded 43 X-ray counterparts within 1 arcmin of the optical galaxy, whereas less than one coincidence is expected by chance. Within this sample of galaxies the vast majority are AGNs (Seyferts 1&2, and LINERS), but there are also some star-forming galaxies and high luminosity early-type galaxies. Further details of the properties of the resulting X-ray sample are given elsewhere (Roberts, 1998). The star-forming galaxies are listed in table 1. In this paper we present X-ray imaging and spectral analysis for the two most luminous star-forming galaxies in the sample: NGC3310 and NGC3690, excluding NGC5905 which underwent some form of transient phenomenon as yet not understood. Both *ROSAT* and *ASCA* observations are available in the HEASARC archive for these two galaxies.

We use a value of $H_0 = 50\text{km s}^{-1}\text{Mpc}^{-1}$ for the Hubble constant, throughout this paper. All errors quoted refer to the 90 per cent confidence level.

1.1 NGC 3310

NGC3310 is a well studied nearby galaxy of Sbc(r)pec type (Mulder et al. 1996). Its recession velocity is 980km s^{-1} (de Vaucouleurs et al. 1991) which implies a distance of 19.6 Mpc. Its optical image shows some interesting features. The most prominent is an arc and bow structure which extends for $100''$ to the North West of the centre of the galaxy. Bertolla and Sharp (1984) propose that the arc could be part of a spiral arm and the bow is the remnant of an old jet. Balick and Heckman (1985) suggest that the whole structure is the remnant of a collision between the galaxy and a

Table 1. The X-ray luminous starburst galaxies in the Ho et al. sample

Galaxy	X – ray luminosity [†]
M82	40.88
NGC3310	40.89
NGC3690	41.62
NGC4449	39.24
NGC5204	39.54
NGC5905	42.18 [‡]

[†] This is the log of the X-ray luminosity in the ROSAT band, from the RASS data.

[‡] This galaxy has show significant long time variability (Bade et al. 1996); This luminosity refers at the outburst epoch.

dwarf companion. This is in agreement with the suggestion of Mulder et al. (1985) who characterize the system as a young merger based on the anomalies found in its rotation curve. It should be noted that in the radio and near-infrared images we do not see two resolved nuclei (Telesco and Gatley 1984, Balick and Heckman 1981). This could mean that the merger phase is now complete as suggested by Balick and Heckman.

Another interesting morphological feature is a giant HII region situated $\sim 12''$ SW of the nucleus (Balick and Heckman 1981). Its size is comparable to the largest extragalactic HII regions known, and its spectral characteristics are dominated by signatures of Wolf Rayet stars (Pastoriza et al. 1993).

The starburst nature of the activity has been confirmed by numerous observations in many spectral bands (e.g. Telesco and Gatley 1984, Smith et al. 1996, Balick and Heckman 1981). Evolutionary synthesis models of the starburst give an estimated age range of 10^7 to 10^8 years, (Van der Kruit and de Bruyn 1976, Balick and Heckman 1981, Telesco and Gatley 1984, Pastoriza et al. 1993).

NGC3310 has been previously observed with the Imaging Proportional Counter (IPC) on board the *EINSTEIN* observatory (Fabianno et al. 1992) yielding an X-ray flux $f_x = 1.1 \times 10^{-12}\text{erg s}^{-1}\text{cm}^{-2}$ in the 0.3-3.5 keV band for a 5keV bremsstrahlung model and Galactic absorption. This corresponds to an X-ray luminosity of $5.0 \times 10^{40}\text{erg s}^{-1}\text{cm}^{-2}$.

1.2 NGC 3690

NGC3690 is also a merging system, but in contrast to NGC3310 it is in an earlier merger phase. The system is composed of NGC3690, which is the western part of the merger and IC694, the eastern component. The whole system is also known as Arp 299 or Mrk 171. Casoli et al. (1989) propose that there is a third component in the system thus forming an interacting triplet. Its recession velocity is 3159km s^{-1} (Sanders and Mirabel 1985), implying a distance of 63.2 Mpc. At that distance the projected separation of $22.5''$ of the two components corresponds to 6.9 kpc. The infrared surface brightness of NGC3690 is at least twice that of IC694 (Friedman et al. 1987). The total mass of gas calculated from radio observations is about $2 \times 10^{11}M_\odot$ (Casoli et al. . 1989)

The high IR luminosity ($L_{FIR} = 1.2 \times 10^{12}L_\odot$) (Soifer et

al. 1987), results from re-radiating dust heated by the starburst activity. This is supported by multi-wavelength studies (Gehrz et al. 1983, Friedman et al. 1987, Nakagawa et al. 1989). Detailed analysis of emission line ratios using the diagnostic diagrams of Veilleux and Osterbrock (1987) clearly indicate a stellar origin of the ionizing continuum (Friedman et al. 1987). The infrared emission is extended over a region of several kpc, but the main sources are located close to the two nuclei (Friedman et al. 1987). Modeling of the starburst has been carried out by Gehrz et al. (1983) and Nakagawa et al. (1989). The latter find a starburst age of about 10 Myrs. An interesting point is that the two nuclei have different properties, implying either a different age and/or a different Initial Mass Function (IMF) for each starburst event.

NGC3690 has been previously observed with the High Resolution Imager (HRI) on board the *EINSTEIN* observatory. 17 counts were detected implying an X-ray flux $f_x = 4.8 \times 10^{-13} \text{ ergs}^{-1} \text{ cm}^{-2}$ in the 0.2-4.0 keV band, for a 5 keV bremsstrahlung model and Galactic absorption (Fabianno et al. 1992). This corresponds to an X-ray luminosity of $2.3 \times 10^{41} \text{ ergs}^{-1} \text{ cm}^{-2}$.

2 OBSERVATIONS AND DATA REDUCTION

2.1 The ROSAT PSPC Observations

NGC3310 and NGC3690 were each observed on two occasions with the Position Sensitive Proportional Counter (PSPC) (Pfefferman et al. 1987) on board *ROSAT* (Trümper et al. 1984). The details of each observation are given in table 2.

For the data reduction we have followed the standard procedure, using the *ASTERIX* package. We excluded those data with Master Veto rates higher than 170 cps. We extracted a PSPC spectral image (channels 11 to 201) with a pixel size of $15''$. To obtain an X-ray spectrum we extracted data from a circular region of $1.5'$ radius around the X-ray centroid. The background was estimated from an annular region between radii of $15'$ and $8.8'$ from the centroid, after exclusion of all the discrete sources detected with the PSS algorithm (Allen 1992) down to the 5σ level.

2.2 The ROSAT HRI Observations

Both galaxies have been observed with the High Resolution Imager (HRI) (David et al. 1997) on board *ROSAT*. HRI covers a field of $38'$ diameter and it has a spatial resolution of about $5''$ (the FWHM of the XRT+HRI Point Spread Function). We note that the HRI has very limited spectral resolution so we cannot use it for spectral analysis (David et al. 1997). The screening of the data has been carried out using the *ASTERIX* package. We have rejected all data with aspect errors greater than 2.

2.3 The ASCA Observations

Both galaxies have been observed with the *ASCA* satellite (Tanaka et al. 1994). On board *ASCA* there are four instruments: two Gas Imaging Spectrometers (GIS2 and GIS3) (Ohashi et al. 1996) and two Solid-State Imaging Spectrometers (SIS0 and SIS1) (Gendreau 1995). For screening of the




Figure 1. The HRI radial profile of NGC3310 (circles) overlaid on the radial profile of AR Lac (stars).

data we have followed the standard procedure and used the ASCACLEAN program in the FTOOLS package with the parameters described in the ABC *ASCA* Guide (Yaqoob et al. 1997). After the standard processing we inspected the light curves and removed time intervals with unusually high numbers of counts, which may result from background particle contamination. The extraction of the source and the background spectra was carried out using the XSELECT program within the FTOOLS package. We extract the source spectrum using a circular region of 2.7 and 5.5 arcmin radius for the SIS and GIS respectively. For the background regions we have selected ten source free zones, each of area about 6 and 10 arcmin² for the SIS and GIS respectively. Following extraction of the source and background spectra, we ran the SISrmg program to create the SIS response matrices for each SIS source spectral file. For the GIS spectral files we used the gis2v4_0.rmf and gis3v4_0.rmf for GIS2 and GIS3 respectively. We created the Ancillary Response Matrices for each file by running the ASCAarf programs within the FTOOLS package.

3 SPATIAL EXTENT ANALYSIS

3.1 NGC3310

We used the archival *HRI* observations to obtain information on the morphology of the X-ray emitting regions and compared these with images of these galaxies at other wavelengths. First we searched for extended emission by comparing the background subtracted radial profile of the galaxies with the radial profile of a point source. For this purpose we used as a reference the point source corresponding to the star AR-Lac. We retrieved archival data for an on-axis observation of AR-Lac, and applied the same extraction process as used for the galaxy. The comparison of the radial profiles of NGC3310, and the radial profile of AR-Lac is shown in figure 1. This figure clearly shows that the X-ray source has a radius of ~ 0.5 arcmin (at larger radii the emission from the galaxy begins to blend with the profile of AR-Lac, within the 2σ error bars). Although this is a somewhat arbitrary point at which to define the extent of the emission, it gives a lower limit to the source size. At the distance of NGC3310 this corresponds to ~ 3 kpc. We note that fitting a Gaussian function to the radial profile gives a similar source size (FWHM=1.6 arcmin).

Table 2. Summary of the observations

Galaxy	Satellite	Instrument	Date Observation started	Exposure Time† Ksec
NGC3310	<i>ROSAT</i>	PSPC	1991-11-16	9.114
	<i>ROSAT</i>	HRI	1995-04-17	41.842
	<i>ROSAT</i>	HRI	1994-11-23	4095
	<i>ASCA</i>		1994-04-17	GIS: 10.4, SIS:10.4
	<i>ASCA</i>		1994-11-13	GIS:11.6, SIS:10.2
NGC3690	<i>ROSAT</i>	PSPC	1993-04-22	3.534
	<i>ROSAT</i>	PSPC	1991-11-18	6.391
	<i>ROSAT</i>	HRI	1993-04-18	6.751
	<i>ASCA</i>		1994-04-06	GIS:5.7, SIS:5.3
	<i>ASCA</i>		1994-12-01	GIS:38.2, SIS:35.5

In order to compare the distribution of the X-ray emitting gas with the optical image we have resampled the X-ray image with a pixel size of 1.5 arc-seconds and then smoothed this image using a two-dimensional gaussian of FWHM=3.5 arc-seconds (2.3 pixels) following Della Ceca et al. (1996). The final pixel size is $5.4''$, equal to the XRT+HRI PSF. Then we obtained contours from this image corresponding to 2.2, 2.0, 1.7, 1.33, 0.88, 0.44, 0.35, 0.26, 0.17 counts arcsec $^{-2}$ and overlaid these onto an optical image obtained from the Digitized Sky Survey. This image is shown in figure 2. The main problem we faced in attempting to overlay the two images is that of frame registration. There are no X-ray sources within the HRI image which have an obvious optical counterpart. Thus in order to overlay the X-ray contour plot onto the optical image, we are forced to assume that the centroid of the X-ray source corresponds to that of the optical nucleus. In the X-ray image of NGC3310 we detect two other sources in addition to the nucleus, at a significance above 10σ . The details concerning these sources are presented in table 3. These point sources may correspond to luminous X-ray binaries or young supernova remnants. Unfortunately there are no identified optical counterparts for these X-ray sources on the POSS plates.

3.2 NGC3690

The HRI analysis of NGC3690 followed the same procedure as for NGC3310. The image pixels were binned to a size of $1.5''$ and then smoothed using a two-dimensional gaussian of FWHM $3.5''$. Finally, we overlaid the X-ray contours on a POSS plate image obtained from the Digitized Sky Survey, figure 3; the contours correspond to 0.57, 0.44, 0.40, 0.35, 0.31, 0.27, 0.22, 0.18, 0.13 and 0.11 counts arcsec $^{-1}$ respectively. The most striking feature is the existence of three separate sources in the X-ray image. The two most luminous sources correspond to the two merging nuclei. Their X-ray fluxes are given in table 4. The third X-ray source corresponds to a hot spot seen in the infrared and radio images to the north of NGC3690 (the western component). Apart from these three sources there are marginal detections of other sources, but at a low level of significance (below 3σ), and will not be discussed further. We note that the relative strengths of the three main X-ray sources follow quite well the relative strengths of the near infrared sources, suggesting a possible common origin for the IR and the X-ray emission.

We have searched for extended emission from the two resolved nuclei. From figure 3 it is clear that the emission

from the two resolved nuclei is extended and not symmetrical. Fitting a Gaussian function to the radial profiles of NGC3690 and IC694 yields a FWHM of 1.5 and 1.8 arcmin respectively.

Comparison between the X-ray images and radio (VLA-A observation of the H92 α radio recombination line, Zhao et al., 1997) and near-IR images (Wynn-Williams et al., 1991), available in the literature, shows some clear similarities. As the radio and near-IR emission is clearly associated with the star-forming regions, the spatial coincidence implies that the soft X-ray emission has its origin in the starburst. We note that we do not see any evidence of X-ray emission from hot connecting the two galaxies, unlike the situation in some other interacting star-forming galaxies (e.g. the Antennae, Fabbiano et al. 1997).

4 SPECTRAL ANALYSIS

We fitted all the datasets simultaneously, thus covering an energy range from 0.1 to 8.0 keV. We rejected all ROSAT PSPC data points below 0.1 and above 2.0 keV; and used only the SIS and GIS data in the range 0.6 to 8.0 keV and from 0.7 to 7.0 keV respectively. In order to apply χ^2 statistics we re-binned the spectra so as to contain at least 20 counts per bin. We used the software package XSPEC (v10) to obtain the spectral fits.

4.1 NGC3310

We first fitted simple one-component models: a thermal bremsstrahlung model, a power-law model and finally a Raymond-Smith plasma model (Raymond & Smith 1977). These models were rejected at greater than the 99.9 per cent confidence level. The results of the spectral fits are given in table 5.

We then fitted two component models: a two temperature Raymond-Smith plasma (ray-ray hereafter) and a Raymond-Smith plasma combined with a power-law (po-ray), following the results of Moran & Lehnert (1997) and Ptak et al. (1997) on the starburst galaxies M82 and NGC253. These models provided a much better fit than the simple models described above, at a confidence level over 90 per cent for an addition of two free parameters (eg Bevington & Robinson 1992). We consider first the po-ray model. We obtained a photon index of $\Gamma = 1.44^{+0.20}_{-0.11}$ and a temperature of $kT = 0.81^{+0.09}_{-0.12}$ keV with a reduced $\chi^2 = 1.115$, where




Figure 2. X-ray contours from the HRI observation of NGC3310 overlaid on a POSS image. The contours are in levels of 2.2, 2.0, 1.7, 1.33, 0.88, 0.44, 0.35, 0.26, 0.17 counts arcsec⁻².

Table 3. Sources in the HRI field of NGC3310

RA (J2000)				Dec (J2000)		Count rate	Flux	Significance
10	38	43.2	+53	31	07	$2.0 \pm 0.27 \times 10^{-3}$	2.3×10^{-14}	15.5
10	38	46.7	+53	30	38	$2.8 \pm 0.26 \times 10^{-3}$	3.07×10^{-14}	23.4

Table 4. Sources in the HRI field of NGC3690

RA (J2000)				Dec (J2000)		Count rate (cps)	Flux (0.1-2.5keV)
11	28	30.7	+58	33	50	$10 \pm 1.2 \times 10^{-3}$	1.18×10^{-13}
11	28	34.1	+58	33	51	$15 \pm 1.5 \times 10^{-3}$	1.72×10^{-13}
11	28	30.0	+58	34	03	$3 \pm 0.6 \times 10^{-3}$	3.17×10^{-14}




Figure 3. X-ray contours from the HRI observation of NGC3690 overlaid on a POSS image. The contours are in levels of 0.57, 0.44, 0.4, 0.35, 0.31, 0.26, 0.22, 0.18, 0.13, 0.11 counts arcsec⁻².

the abundance is fixed to the solar value and the absorbing column density to the Galactic value, $N_H = 0.7 \times 10^{20} \text{ cm}^{-2}$, (Stark et al. 1992). Leaving the absorbing column density as a free parameter further improves the fit (reduced $\chi^2 = 1.031$); this is significant at a confidence level of > 99 per cent for one additional parameter. In contrast, leaving the abundance as a free parameter does not improve the fit further at a statistically significant level.

In the ray-ray case we do not obtain a significantly improved fit compared to the po-ray model (reduced $\chi^2 = 1.022$). Figure 4 shows the spectrum of NGC3310 with the double Raymond-Smith model. We obtained temperatures of $0.80^{+0.07}_{-0.04} \text{ keV}$ and $14.98^{+13.52}_{-4.88} \text{ keV}$ for an absorbing hydrogen column density of $N_H = 1.37^{+0.50}_{-0.32} \times 10^{-20} \text{ cm}^{-2}$. When we fixed the column density to the Galactic value ($N_H \approx 1.0 \times 10^{20} \text{ cm}^{-2}$) we obtained $0.84^{+0.05}_{-0.07} \text{ keV}$ and $kT \approx 64 \text{ keV}$. We also introduced two different absorbing column densities, for the soft and the hard components, following the

analysis of M82 by Moran & Lehnert (1997) who showed that the hard X-ray component is obscured at the nucleus by $N_H \sim 10^{22} \text{ cm}^{-2}$. However, we do not find similar evidence here, since the hard component column has a value of $N_H \approx 1.77 \times 10^{20} \text{ cm}^{-2}$. Finally, we attempted to measure the element abundance in order to compare it with that for other star-forming galaxies (Sansom et al. 1996, Serlemitsos et al. 1996), which appear to show a systematic trend towards sub-solar abundances. Unfortunately the data for our galaxies are not able to set any useful constraint on the abundances.

We do not detect the $FeK\alpha$ line at 6.7 keV as would be expected for a hot thermal plasma. The low signal/noise data above $\sim 5 \text{ keV}$, does not allow us to set a useful upper limit on the line equivalent width. The lack of a strong FeK line could be explained by lower than solar abundances, as is probably the case in other well-studied starbursts such as




Figure 4. The top panel shows *ROSAT* and *ASCA* spectra of NGC3310 with the best fit double Raymond Smith model. The bottom panel shows the ratio of the data points to the model.

The SIS data are marked with dots while the PSPC data are marked with triangles.

NGC253, (Ptak et al. 1997), and the Antennae, (Sansom et al. 1996).

We note that a previous preliminary analysis of *ASCA* data by Serlemitsos et al. (1996), find a spewctral slope consistent with our value. However, they derive a considerably higher column density, but without the benefit of using the lower energy *ROSAT* data.

4.2 NGC3690

The spectral analysis follows the same procedures as for NGC3310. Simple one-component models such as a power-law and a Raymond-Smith plasma are similarly rejected (see table 6). Two-component models, such as a po-ray and a ray-ray model (with the hydrogen column density as a free component) give $\Gamma = 1.56^{+0.11}_{-0.11}$, $kT = 0.83^{+0.02}_{-0.04}$, $N_H = 2.4^{+0.6}_{-0.5} \times 10^{20} \text{ cm}^{-2}$ and $kT_1 = 0.83^{+0.03}_{-0.03}$, $kT_2 = 10.3^{+5.95}_{-2.44}$, $N_H = 1.6^{+0.4}_{-0.4} \times 10^{20} \text{ cm}^{-2}$ respectively (the Galactic absorption is $N_H = 1 \times 10^{20} \text{ cm}^{-2}$). In the above model the abundance was fixed to solar. Also, as for NGC3310 when the abundance is a free parameter, it cannot be usefully constrained. Figure 5 shows the *ROSAT* and *ASCA* spectrum of NGC3690 with the best fit double temperature Raymond-Smith thermal plasma model. We can clearly see from the spectrum that there is no evidence for an Fe line at ~ 6.7 keV. The upper limit to the Fe line equivalent width is $\sim 860\text{eV}$. We also tried to fit the models using a different absorbing column for the hard component. We obtain $N_H = 1.7^{+0.1}_{-0.8} \times 10^{20} \text{ cm}^{-2}$, which is comparable to the previous best fit values, while the χ^2 is not significantly improved for one additional parameter. Thus there is no evidence for excess absorption in the hard component, in contrast to the results for M82 (Lehnert & Moran 1997). We also note that both the ray-ray and the po-ray models do not provide an adequate fit to the data as they can be rejected at over the 99 per cent level of confidence using a χ^2 goodness of fit.

We therefore attempted to fit more complicated models, such as multi-temperature components, and a power-law distribution of temperatures. This is model ‘cevmecl’ in XSPEC, (see Done and Osborne 1997). This model gives a




Figure 5. The top panel shows *ROSAT* and *ASCA* spectra of NGC3690 with the best fit double Raymond Smith model. The bottom panel shows the ratio of the data points to the model.

The SIS data are marked with dots, while the PSPC data are marked with triangles.

best fit of reduced $\chi^2 = 1.286$, with a maximum temperature of ~ 57 keV, a power-law index of the temperature distribution of $\alpha = 0.22^{+0.11}_{-0.10}$ and $N_H = 3.21^{+0.73}_{-0.48} \times 10^{20} \text{ cm}^{-2}$. A combination of a power-law and the multi-temperature model gives a slightly better fit (reduced $\chi^2 = 1.211$) with $\Gamma = 1.52^{+0.13}_{-0.12}$, $T_{\text{max}} = 1.17^{+0.25}_{-0.16}$ and α fixed to 1. Finally, we considered using a non-equilibrium-ionization model (NEI), such as that applied to hot gas resulting from supernova explosions (Hughes and Singh 1994). This model assumes that the gas is instantaneously shock heated, and then applies corrections to the Raymond-Smith thermal plasma code to account for the non-equilibrium ionization fraction. An additional variable in the NEI model is nt , which is the product of the electron density and the time elapsed since the passage of the shock wave. In principle the results from fitting NEI models can be used to estimate the characteristic timescale over which the gas reaches ionization equilibrium. Clearly for a situation in which multiple supernova occur over an extended time period, it will not be possible to identify a unique timescale. However, NEI models have been applied to the optically thin plasma in galactic scale superwinds, such as observed in M82, Tsuru et al (1998). Unfortunately in the case of NGC3690, we do not have the benefit of measurements of the low energy lines of silicon and magnesium needed to quantify the importance of non-equilibrium effects. For NGC3690 the NEI models give a best fit with a combination of a power-law and a NEI model. However the reduction in the χ^2 is not statistically significant at a level above 90%, and so we conclude that our data does not justify further consideration of the NEI models. However, if new soft X-ray emission line data become available, it would be worthwhile returning to this question.

5 DISCUSSION

5.1 The soft X-ray emission

The HRI images clearly show the extended X-ray emission in both NGC3310 and NGC3690/IC694. This supports a thermal origin for the soft emission, which may be due to

Table 5. Spectral fitting results for NGC 3310

Parameter	Power-law	Single Temperature Raymond-Smith	Double Temperature Raymond Smith	Raymond Smith + Power-law
kT (KeV)		$7.35^{+1.24}_{-1.18}$	$0.80^{+0.07}_{-0.04}$ $14.98^{+13.52}_{-4.88}$	$0.81^{+0.09}_{-0.12}$
Γ	$1.68^{+0.08}_{-0.08}$			$1.44^{+0.20}_{-0.11}$
$N_H(10^{20} \text{ cm}^{-2})$	$3.15^{+0.63}_{-0.52}$	$1.90^{+0.45}_{-0.36}$	$1.37^{+0.50}_{-0.32}$	$1.74^{+0.68}_{-0.40}$
$\chi^2 / \text{d.o.f.}$	222.9/175	239.3/175	168.7/165	170.2/165
Flux † (0.1-2.0keV)	0.95	0.86	0.9	1.09
Flux † (2.0-10.0keV)	1.88	2.12	2.10	2.10
Total Luminosity ‡ (0.1-10keV)	1.3	1.4	1.4	1.5

† All the fluxes are in units of $10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$. ‡ The luminosity is in units of $10^{41} \text{ rmerg s}^{-1}$.

a galactic scale super-wind eg Heckman et al. (1993). Such a phenomenon is observed in other star-forming galaxies, M82 (Strickland, Ponman & Stevens 1997), NGC1569 (Della Ceca et al. 1996) and NGC4449 (Della Ceca et al. 1997). According to this scenario, supernova in the starburst region create a hot ($\sim 10^8$ K) bubble. The dense shell around the bubble fragments due to Rayleigh-Taylor instabilities and forms the optical emission line filaments and is probably a source of soft X-ray emission. The optical data also support this model. In the case of NGC3690 the H_α images show plumes and filaments extending out to several kpc (Armus, Heckman & Miley 1990). Additional evidence comes from the extended synchrotron radio emission which may arise from the electrons produced by supernova remnants in the starburst regions, (Gehrz et al. 1983). In the case of NGC3310 the situation is more complicated as its low inclination angle ($\theta \approx 32^\circ$) makes detection of any outflow along the galaxy's minor axis very difficult.

In order to further test the validity of the above model, we can estimate the expected X-ray luminosity of the super-wind and compare this with the observed luminosity of the soft Raymond-Smith component. First we relate the X-ray luminosity of the gas contained in a superbubble with the basic properties of the starburst; its bolometric luminosity and age. Heckman et al. (1996) have calculated the total X-ray luminosity from a superbubble by integrating over its volume, and using the expressions for the gas density and the bubble radius derived by MacLow and McCray (1988). They find that $L_X \simeq 5 \times 10^{39} L_{41}^{33/35} n^{17/35} t_7^{19/35} \text{ erg s}^{-1}$, where L_{41} is the injection rate of mechanical energy into the bubble in units of $10^{41} \text{ erg s}^{-1}$, n is the gas density and t_7 is the age of the bubble in units of 10Myrs. Considering that the formation of the bubble begins, as a result of strong winds from O stars following the starburst, we can assume that the age of the bubble is almost the same as the age of the starburst. Using the relation $dE/dt = 7 \times 10^{42} L_{\text{IR},11} \text{ erg s}^{-1}$, (Heckman et al. 1996) which correlates the total infrared luminosity (in units of $10^{11} L_\odot$) of a starburst with age greater than 10^7 yr with the kinetic energy deposition rate from supernovae and stellar winds, we can estimate the X-ray luminosity. Using the IRAS data from Soifer et al. (1987), the IR luminosity of NGC3310 is $5.5 \times 10^{10} L_\odot$. Assuming a gas density of $\sim 1 \text{ cm}^{-3}$ and a typical starburst age of 10Myr we obtain an X-ray luminosity $L_x = 1.5 \times 10^{41} \text{ erg s}^{-1}$ in the ROSAT band. This luminosity is larger than the measured luminosity in the same energy band of ($L_X = 4.7 \times 10^{40} \text{ erg s}^{-1}$), but considering the large uncertainties in the model parameters involved, we conclude that the super-wind model remains a

possible scenario for the origin of the soft X-ray emission. In NGC3690/IC694 the situation is more complex, as there are at least three distinct starburst components (Nakagawa et al. 1989) which are unresolved by IRAS. In order to obtain an estimate of the mechanical energy input for each component, we used the ground based mid-IR data (10-32 microns) of Wynn-Williams et al. (1991). These observations have sufficient spatial resolution to separate the three components. We then make the reasonable assumption that the fractions of the total IRAS flux which arise from each of the three star-forming components, are similar to the fractions in the mid-IR band. Using the same formulae (Heckman et al. 1996) as in the analysis of NGC3310 above, and assuming a starburst age of 10 Myr and a density of $n = 1 \text{ cm}^{-3}$, we predict $L_x = 1.3 \times 10^{42}$, 1.1×10^{42} and $3.2 \times 10^{41} \text{ erg s}^{-1}$ for the A,B and C components respectively (following notation of Gehrz et al., 1983). Again the predicted luminosity is higher than the measured soft (0.1-2.5keV) X-ray luminosity ($4.7 \times 10^{41} \text{ erg s}^{-1}$). This could possibly result from the fact that some fraction of the far-infrared emission measured by IRAS, may arise from regions more extended than those producing the X-rays. Another possibility is that an additional source of heating may be present, perhaps due to stars of later spectral type than those producing the winds.

The soft X-ray spectra of NGC3310 and NGC3690 are very similar. A Raymond-Smith component with a temperature of kT ~ 0.8 keV dominates the spectrum at soft energies and could originate from the super-wind. This temperature is similar to that found for other star-forming galaxies, eg M82 and NGC253 (Ptak et al. 1997), NGC1569 (Della Ceca et al. 1996), NGC4449 (Della Ceca et al. 1997) NGC6240 and NGC2782 (Schulz et al. 1997). In the case of NGC3690 where the photon statistics are good, the SIS spectral fit yields a poor χ^2 fit. This could be suggestive of a hot tenuous gas component. Indeed, if the expanding gas has not had time to reach thermal equilibrium, its spectrum will not be well represented by the Raymond-Smith model. The use of a non-equilibrium code by Hughes et al. improved the fit, but again the χ^2 is not accepted at the 98 per cent confidence level. This is not surprising as even a non-equilibrium model may be inadequate to fit an ensemble of supernova remnants occurring in several star-forming regions as is the case for NGC3690/IC694.

5.2 The hard X-ray emission

The spectrum at X-ray hard energies can be represented, in both galaxies, by a power-law ($\Gamma \sim 1.4 - 1.6$), or alter-

natively by a Raymond-Smith component with high temperature, ($kT > 10$ keV). There is no strong evidence for the presence of a large amount of obscuration as is the case for NGC253 and M82 (Ptak et al. 1997). There are at least four possibilities for the origin of the hard X-ray emission: a) Inverse Compton scattering of the IR photons produced in the starburst by electrons from the numerous supernova remnants (eg Schaaf et al. 1989) b) emission from a low-luminosity AGN as is the case in NGC3628 (Yaqoob et al. 1995) c) thermal emission from a very hot gas ($T \sim 10^8$ K) and d) emission from X-ray binaries (Griffiths & Padovani 1990). We discuss each of these possibility in turn. We consider first the possibility, that the hard X-ray emission arises from Inverse Compton (IC) scattering of the copious infrared photons off of the relativistic electrons generated by the numerous supernovae. Support for this, especially in the case of NGC3310, comes from the similarity (within the errors) of the spectral index of the hard X-rays and the spectral index of the radio emission ($\alpha_{rad} = 0.61^{+0.03}_{-0.03}$ Niklas et al. 1997). Following Schaaf et al. (1989) we can estimate the X-ray luminosity from Inverse Compton scattering. Unfortunately, we can perform this calculation only for NGC3310, as there are no suitable radio data available for NGC3690. From Vallée (1993) we have that the “minimum energy” magnetic field (see Longair 1992) is $B = 0.47 \times 10^{-5}$ Gauss. Then the minimum energy density is $u = (7/3)(B^2/8\pi) \text{ erg cm}^{-3} = 2.05 \times 10^{-12} \text{ erg cm}^{-3}$. Following Schaaf et al. (1989) we have that $L_{IC} = (1/3)\sigma_T R_{IC} L_{IR} \frac{\epsilon_e}{mc^2} \gamma_2^{0.8} \gamma_1^{0.2}$, where σ_T , R_{IC} , L_{IR} and mc^2 , ϵ_e are the Thomson cross-section, the thickness of the disk, the far-IR luminosity, the rest mass of the electron and the energy density of the relativistic electrons, respectively. For a typical galactic disk $R_{IC} \sim 1 \text{ kpc}$ (we cannot measure the actual thickness of the disk since NGC3310 is almost face on) and for a typical value of the low frequency cut-off $\nu = 0.01 \text{ GHz}$ we have $\gamma_1 = 150$ for the lower limit of the Lorentz factor, and $\gamma_2 = 10^3$ for the maximum Lorentz factor, in order to have IC emission at 10keV (Schaaf et al. 1989). So for the Inverse Compton X-ray luminosity we predict a value of $2.5 \times 10^{38} \text{ erg s}^{-1}$, much lower than the detected hard X-ray luminosity of NGC3310, thus implying that IC can be only a minor component of the total X-ray emission from this galaxy. We caution that this result is quiet uncertain as the calculation depends on parameters like the volume of the source and the thickness of the disk which are poorly known. Another possible origin of the hard X-ray emission is the presence of a low luminosity AGN, although there is no evidence for non-starforming nuclear activity from diagnostic emission line ratio diagrams based on optical and near infrared spectra.

However, the hard X-ray power-law spectrum is nevertheless consistent with the presence of an active nucleus. In order to test this possibility further we use the L_X/L_{H_α} relation from Elvis et al. (1984), where L_{H_α} is the luminosity of the broad component of the H_α line and L_X is the hard X-ray luminosity (2-10 keV). $L_X/L_{H_\alpha} \simeq 40$, for low-luminosity AGN and thus we estimate $f(H_\alpha) \sim 5.3 \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$ and $2.7 \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$ for NGC3310 and NGC3690 respectively. A broad component of this strength is easily detectable, but is not seen (Ho et al. 1997). The absorbing columns found from the X-ray spectral fitting (ray-po models in tables 5 and 6) imply low

extinctions, $A_V \sim 0.1$ mag for both galaxies (Bohlin et al. 1978). Hence a BLR reddened by this amount would still be observed.

The next possibly is the presence of a very hot thermal component. A Raymond-Smith model with $kT > 10$ keV provides a good fit to the data. However, the strong FeK line at 6.7 keV which should accompany the thermal emission is not observed. This could be attributed to a low metallicity, but unfortunately we cannot check this as the abundances are not well constrained using the present X-ray data. Alternatively, the lack of a FeK line could be explained by a low contribution of type Ia supernovae to the enrichment of the Interstellar Medium (Sansom et al. 1996).

Finally we consider high mass X-ray binaries as a possible origin of the hard X-ray component. High mass binaries will form as a consequence of the starburst, and indeed many such systems have been identified in nearby star-forming galaxies (Read et al. 1997, Fabbiano 1995 and references therein). Assuming that a typical X-ray luminosity of these systems is $10^{37-38} \text{ erg s}^{-1}$, we can estimate the total number of binaries. As the X-ray luminosity of the hard component is $L_x \sim 5 \times 10^{41}$ and $L_x \sim 10^{41} \text{ erg s}^{-1}$ for NGC3690 and NGC3310 respectively, we estimate a range of between 5,000-50,000 and 1,000-10,000 X-ray binaries for the two galaxies. Now we can compare this with the number of ionizing OB stars determined from the integrated far-infrared luminosity. Making the assumption that it is mostly these stars that heat the dust, which then reradiates in the mid-far infrared, we estimate $\sim 2 \times 10^5$ and $\sim 3.5 \times 10^6$ OB stars for NGC3310 and NGC3690, respectively. If, following Fabbiano et al. (1992), 0.2 percent of these are massive X-ray binaries, then there are 400, and 7000 such systems, respectively. For the upper range of binary luminosities the predicted X-ray luminosity is comparable to that observed in both cases. Indeed, if the hard X-ray emission arises from binaries with low-metallicity and thus high X-ray luminosity ($L_x \sim 10^{38-39} \text{ erg s}^{-1}$) like those observed in the Magellanic Clouds (van Paradijs & McClintock 1995), then they could easily produce the observed luminosity. One potential problem is that some point sources observed in nearby galaxies with the ROSAT PSPC by Read et al. (1997), appear to have soft spectra ($kT \sim 2$ keV), whilst the the high temperatures inferred for NGC3310 and NGC3690 are closer to those predicted for high mass X-ray binaries (see Nagase 1989).

6 CONCLUSIONS

We have modelled and interpreted the combined ASCA and ROSAT X-ray spectra from 0.1-10 keV, for the star-forming galaxies NGC3310 and NGC3690. These two galaxies are amongst the most X-ray luminous ($L_x \sim 10^{41-42} \text{ erg s}^{-1}$) starbursts in the local Universe. In addition, ROSAT HRI images show that the emission from NGC3310 is extended out to at least ~ 3 kpc. The soft X-ray emission from NGC3690 comes from at least three spatially resolved regions. The limited spatial resolution of the ASCA data does not allow us to place useful limits on the extent of their hard X-ray components. The X-ray spectrum of NGC3310 can be described by two components: at soft energies a Raymond-Smith component ($kT \sim 0.8$ keV) which probably originates

from a super-wind. The predicted soft X-ray emission, on the basis of the supernovae mechanical energy deposited to the interstellar medium, is comparable to that observed. At harder energies we can fit either a Raymond-Smith component ($kT \sim 17$ keV) or a power-law $\Gamma \sim 1.4$.

The results for NGC3690 are similar. There are at least two components in its spectrum: a soft Raymond-Smith plasma ($kT \sim 0.8$ keV) and a harder component which can be represented equally well by a Raymond-Smith component ($kT \sim 10$ keV) or a hard power-law $\Gamma \sim 1.6$. However, the best-fit model for NGC3690 is rejected at the 98 per cent confidence level implying that more complicated models are necessary. Hence we considered a non-equilibrium ionization model (Hughes et al. 1994) and a multi-temperature thermal model (Done & Osborne 1996). The above models do improve the fit, but only at the ~ 90 per cent confidence level. Although non-equilibrium models may be important, additional information such as X-ray emission line ratios are required in order to make further quantitative progress.

In neither NGC3310 and NGC3690 do we find evidence for significant absorption columns above the Galactic values, in either the soft or the hard X-ray component.

The nature of the hard X-ray emission is still uncertain, and we considered various possibilities. Of these we conclude that Inverse Compton scattering by high energy radio electrons and infrared photons, can probably provide only a minor contribution. Although the presence of an AGN is a possibility, this is not supported by data at other wavelengths, and the modest columns argue against an obscured nucleus. A very hot thermal component may be present, ($\sim 10^8$ K). We do not detect the FeK line at 6.7 keV, possibly because the abundances are sub-solar. To test this possibility further requires information on element abundances, and an extension of the observed spectra to higher energies. Finally, estimates of the number of X-ray binaries, based on the total mid-IR emission, suggests that they may well be able to account for the hard X-ray emission.

The X-ray results on these two galaxies are similar to those obtained for other dwarf star-forming galaxies (eg NGC1569, NGC4449) and the archetypal star-forming galaxies M82 and NGC253, although there is a range in relative contributions from the various X-ray components. This implies that the same general emission mechanisms apply in star-forming galaxies over three decades of luminosity.

7 ACKNOWLEDGMENTS

We are grateful to J. Hughes for providing the supernova remnant non-equilibrium XSPEC model.

REFERENCES

Allen D.J., ASTERIX User Note 004, STARLINK, 1992
 Armus L., Heckman T.M., Miley G.K., 1990, ApJ, 364, 471
 Bade N., Komossa S. and Dahlem M., 1996, A&A, 309, 35L
 Balick B. and Heckman T., 1981, A&A, 96, 271
 Bertolla A.F., and Sharp N.A., 1984, MNRAS, 207, 47
 Bevington P.R. and Robinson D.K., 1992, Data reduction and error analysis for the physical sciences, 2nd ed., McGraw Hill

Bohlin R.C., Savage B.D. and Drake J.F., 1978, ApJ., 224, 132
 Casoli F., Combes F., Augarde R., Figon P. and Martin J.M., 1989, A&A, 224, 31
 David et al. 1997 The *ROSAT* HRI Calibration Report
 hea-www.harvard.edu/rosat/rsdc_www/HRI_CAL_REPORT
 de Vacouleurs G. et al. , 1991, Third Reference catalogue of bright galaxies.
 Dekel A. and Silk J., 1986, ApJ, 303, 39
 Della Ceca R., Griffiths R.E., Heckman T.M. and Mackenty J.W., 1996, ApJ, 469, 662
 Della Ceca R., Griffiths R.E. and Heckman T.M., 1997, ApJ, 485, 581
 Done C.D. and Osborne J.P., 1997, MNRAS, 288, 649
 Elvis M., Soltan A. and Keel W., 1984, ApJ., 283, 479
 Fabbiano G., 1988, ApJ, 3310, 672
 Fabbiano G., 1995, in X-ray Binaries edited by Lewin W.H., van Paradjis J., and van den Heuvel E.P.J.
 Fabbiano G. and Trinchieri G., 1984, ApJ, 286, 491
 Fabbiano G., Kim D.W., and Trinchieri G., 1992, ApJS, 80, 531
 Fabbiano G., Schweizer F. and Mackie G., 1997, 478, 542
 Friedman S.D., Cohen R.D., Jones B., Smith H.E. and Stein W.A., 1987, AJ, 94, 1480
 Gehrz R.D., Sramek R.A., Weedman D.W., 1983, ApJ, 267, 551
 Gendreau K.C., 1995, PhD. Thesis
 Griffiths R.E. and Padovani P., 1990, ApJ., 360, 483
 Heckman T.M, Armus L., Miley G.K., 1990, ApJS, 74, 833
 Heckman T.M, Dahlem M, Lehnert M.D., Fabbiano G., Gilmore D., Waller W.H., 1995, ApJ., 448, 98
 Ho L.C., Filippenko A.V., Sargent W.L., 1995, ApJS, 98, 477
 Ho L.C., Filippenko A.V., Sargent W.L.W., 1997, APJS, 112, 315
 Hughes J.P. and Singh K.P., 1994, ApJ, 422, 126
 Kim D.W., Fabbiano G., Trinchieri G., 1992, ApJS, 80, 645
 Lehnert M.D. and Heckman T.M., 1996, ApJ, 462, 651
 Long K.S. and van Spreybroeck L.P., 1983, in Accretion driven stellar X-ray sources, edited by Lewin W.H.G. and van Den Heuvel E.P.J., CUP
 Longair M.S. 1992, High Energy Astrophysics vol.2, CUP
 MacLow M.M., and McCray R., 1988, ApJ, 324, 776
 Moran E.C., Halpern J.P., Helfand D.J., 1996, ApJS, 106, 341
 Moran E.C. and Lehnert M.D., 1997, ApJ, 478, 172
 Mulder P.S. and Van Driel W., 1996, A&A, 309, 403
 Nagase F., 1989, PASJ, 41, 1
 Nakagawa T., Nagata T., Geballe T.R., Okuda H., Shibai H., Matsuhara H., 1989, ApJ, 340, 729
 Niklas S., Klein U., Weilebinski R., 1997, A&A, 322, 19
 Ohashi et al. , 1990, ApJ, 365, 180
 Ohashi et al. , 1996, PASJ, 48, 157
 Pastoriza M.G., Dottori H.A., Terlevich E., Terlevich R., Diaz A.I., 1993, MNRAS, 260, 177
 Pfefferman et al. , 1986, Proc. SPIE, 733, 519
 Ptak A., Serlemitsos P., Yaqoob T., Mushotzky R., Tsuru T., 1997, AJ, 113, 1286
 Raymond J.C., Smith B.W., 1977, ApJSS, 35, 419
 Read A.M., Ponman T.J., Strickland D.K., 1997, MNRAS, 286, 626

- Read A.M., Ponman T.J., Wolstencroft R.D., 1995, MNRAS, 277, 397
- Rieke G.H., Lebofsky M.J., Thompson R.I., Low F.J., Tokunaga A.T., 1980, ApJ, 238, 24
- Roberts T., 1998, PhD Thesis, University of Leicester, in preparation
- Sanders D.B. and Mirabel I.F., 1985, ApJ, 298, 31L
- Sansom A.E., Dotani T., Okada K., Yamashita A., Fabbiano G., 1996, MNRAS, 281, 48
- Schaaf R. et al. , 1989, ApJ, 336, 722
- Schultz H. et al. , 1997, preprint
- Serlemitsos P., Ptak A., Yaqoob T., 1996, in The physics of LINERS in view of recent observations, 1996, edited by M. Eracleous, A. Koratkar, C. Leitherer, and L. Ho
- Smith et al. ,1996, ApJ, 473, L21
- Soifer B.T. et al. , 1987, ApJ, 320, 238
- Sutherland R.S., Dopita M.A., 1993, ApJS, 88, 253
- Stark A. A. et al. , 1992, ApJS, 79, 77
- Stewart G.C., Fabian A.C., Terlevich R.J., Hazard C., 1982, MNRAS, 200, 61
- Strickland D.K., Ponman T.J., Stevens I.R., 1997, A&A, 320, 378
- Tanaka Y., Inoue H., Holt S.S., 1994, PASJ, 46, 37L
- Telesco C.M. and Gatley I., 1984, ApJ, 284, 557
- Trümper J., 1984, Physica Scripta, T7, 209
- Tsuru T.G., Awaki H., Koyama K., Ptak A., 1998, PASJ, 49, 619
- Vallée J.P., 1993, MNRAS, 264, 665
- Van der Kruit P.C. and de Bruyn A.G., 1976, 48, 373
- Van Paradjis J. and McClintock J.E., 1995, in X-ray Binaries edited by Lewin W.H., van Paradjis J., and van den Heuvel E.P.J.
- Veilleux S., Osterbrock D.E., 1987, ApJS, 63, 295
- Voges W. et al. , 1996, IAU Circ. 6420
- Watson M.G., Stanger V., Griffiths R.E., 1984, ApJ, 286, 144
- Wilson A.S. et al. , 1992, ApJ, 391, L75
- Wynn-Williams et al. ., 1991, ApJ, 377, 426
- Yaqoob T. et al. , 1995, ApJ, 455, 508
- Yaqoob T. et al. , 1997, The ASCA ABC Guide, v2.0, NASA/GSFC
- Zhao J.H., Anantharamaiah K.R., Goss W.M., Viallefond F., 1997, ApJ, 482, 186